



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/21	A1	(11) International Publication Number: WO 95/24212 (43) International Publication Date: 14 September 1995 (14.09.95)
(21) International Application Number: PCT/GB95/00488 (22) International Filing Date: 7 March 1995 (07.03.95) (30) Priority Data: 9404379.1 7 March 1994 (07.03.94) GB 9420340.3 10 October 1994 (10.10.94) GB (71) Applicant (for all designated States except US): IMPERIAL COLLEGE OF SCIENCE, TECHNOLOGY & MEDICINE [GB/GB]; Sherfield Building, Imperial College, London SW7 2AZ (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): FOSTER, Graham, Russell [GB/GB]; Departement of Medicine, Queen Elizabeth the Queen Mother Wing, St Mary's Hospital Medical School, London W2 1PG (GB). THOMAS, Howard, Christopher [GB/GB]; Department of Medicine, Queen Elizabeth the Queen Mother Wing, St Mary's Hospital Medical School, London W2 1PG (GB). (74) Agents: CHAPMAN, Paul, William et al.; Kilburn & Strode, 30 John Street, London WC1N 2DD (GB).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published <i>With international search report.</i>
(54) Title: THE USE OF INTERFERON SUBTYPES IN THE PREPARATION OF MEDICAMENTS TO TREAT VIRAL INFECTIONS		
(57) Abstract Individual subtypes of interferon- α (IFN- α) have been found to have different antiviral activity in different cell types and are therefore used to prevent or treat viral infections in cell types in which they are most active. The individual subtype of choice has relatively low antiviral activity in other celltypes to reduce the risk of side effects. IFN- α_{10} and IFN- α_{17} are preferred for use in treating viral lung infections and IFN- α_8 is preferred for use in treating viral liver infections.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

The Use of Interferon Subtypes in the Preparation of
Medicaments to Treat Viral Infections

5 This invention relates to the treatment or prophylaxis of
viral infections with interferons (IFNs).

10 Type I interferons (IFN) are a family of closely related
glycoproteins containing many IFN- α subtypes and one IFN- β
subspecies. At least 23 different human IFN- α subtypes
have been identified by analysis of human cDNA libraries
and by protein analysis of the IFNs produced by
stimulated lymphoblastoid cells. The reasons for this
heterogeneity are not yet known. Previous studies have
suggested that all Type I IFNs bind to an identical
15 receptor and therefore have identical effects. However
a mutant cell line that responds only to IFN- β but not
IFN- α has been identified showing that these two IFN
subspecies bind to a different receptor and may therefore
have different effects. Studies on the recently
20 identified transmembrane human IFN receptor have shown
that if this receptor is transfected into murine cells
the cells respond only to some IFN subtypes, showing that
a second receptor component is required to allow cells to
respond to IFN and that the murine equivalent of this
25 component is able to distinguish between different IFN
subtypes. Molecular analysis of the human Type I IFN
receptor thus suggests that the receptor may be able to
distinguish between different IFN subtypes, but whether
the different subtypes do, in fact, have different
30 effects is not yet clear. A number of studies have
compared the effects of different IFN- α subtypes on the
antiviral activities of human cell lines. Zoon et al (J.
Biol. Chem. 267: 15210-16 (1992) studied IFNs that were
obtained from HPLC purification of natural IFN and found

no gross differences in their antiviral activities. However, Sperber et al, *J. Interferon. Res.* 12 363-368 (1992) examined the effects of different recombinant IFN- α subtypes on cells infected with the human immunodeficiency virus (HIV) and found marked differences in their antiviral properties.

Whereas the investigations of Sperber et al were confined to the effect of different subtypes of IFN- α against a particular virus (HIV-1), it has now been found that the antiviral effect of subtypes of IFN- α is dependent on the type of cell infected with the virus. Further, it appears that certain subtypes of IFN- α act as partial agonists to antivirally effective IFNs- α . Therefore, by virtue of the invention, the use of specific subtypes of IFN- α for the treatment of each cell type is indicated.

According to the invention, there is provided the use of a single interferon- α (IFN- α) subtype in the preparation of a medicament for preventing or treating viral infections of a particular organ or cell type.

The cell type will generally not be T-lymphocytes, in view of the prior work of Sperber et al. However, nothing in the Sperber et al publication referred to above suggests that IFNs- α exhibit cell-type specific antiviral activity.

A particularly preferred IFN- α subtype is IFN- α_8 . This is particularly suitable for treating or preventing viral infections of the liver. In addition to its potent antiviral effects in normal cell lines, IFN- α_8 is also active in a mutant cell line (11,1 (Pellegrini et al, *Mol. Cell. Biol.* 9: 4605-4612 (1989))) that does not respond to

other α IFN subtypes.

The particular IFN- α subtype to be used in clinical practice will depend on the cell type which is infected.
5 Preferred subtypes for a particular cell type may be those which not only have potent antiviral activity for that particular cell type but also have relatively low activity in respect of other cell types, so as to reduce the possibility of side effects.

10 For example, when choosing an IFN- α subtype for use in lung infections, regard will be had to *in vitro* studies on lung carcinoma cell lines which showed that IFN- α_2 , IFN- α_5 , IFN- α_8 , IFN- α_{14} and IFN- α_{17} were the most potent
15 subtypes tested and IFN- α_{10} had high activity. However, IFN- α_2 , IFN- α_5 , IFN- α_8 and IFN- α_{14} were also very potent antivirals in liver cell lines. Thus, it may be the case that the preferred subtypes for treating or preventing viral infections of the lung are IFN- α_{10} and IFN- α_{17} .

20 When choosing an IFN- α subtype for use in liver infections, regard will also be had to *in vitro* studies on liver cell lines which showed that IFN- α_2 , IFN- α_5 and IFN- α_8 were the most potent subtypes tested. IFN- α_2 and
25 IFN- α_5 were, as mentioned above, also potent antivirals in a liver cell line. However, although this is also true of IFN- α_8 , it has also been observed that, *in vitro*, cells appear to produce IFN- α_8 in response to viral infection. Thus, the preferred subtype for treating or preventing
30 viral infections of the liver is IFN- α_8 .

Mixtures of a small number (such as two, three or four) specific IFNs- α are also contemplated within the scope of the invention. Each will generally be selected in

accordance with the guidelines given above.

IFN- α subtypes may be administered by conventional means and at doses comparable to those known in the art, although the precise mode of administration and dosage will always be within the discretion of the physician or other medical practitioner.

The invention has application in a method of preventing or treating viral infections of a particular organ or cell type, the method comprising administering an effective antiviral amount of a single interferon- α (IFN- α) subtype.

The invention will now be illustrated by the following example.

The example refers to Figure 1 in which:

FIGURE 1: shows the relative ED 50 for the various IFN- α subtypes against various cell lines.

EXAMPLE

25

a) Interferons

WELLFERON[®] lymphoblastoid interferons were obtained from The Wellcome Foundation Limited. Human Type I interferons were prepared by stimulating Namalwa cells with Sendai virus and then purifying and fractionating the IFN mixture using antibody precipitation and HPLC purification, as previously described (Zoon et al, *J. Biol. Chem.* 267: 15210-15216 (1992)). IFNs were also prepared, in a similar manner, from supernatants of Sendai virus-treated human peripheral blood mononuclear

cells (Interferon Sciences Inc). The identity of the purified IFN subtype fractions was confirmed by microsequencing a fraction of the column eluate and the concentration of the final product was determined using a commercial kit (Sigma).

b) Antiviral assays

Human cell lines (HuH7, A549 and SHSY) were grown in DMEM supplemented with 10% FCS. Antiviral assays were performed as described (Zoon et al, supra). In brief, cells were transferred to 96 well microtitre plates (1.5×10^4 cells per well) and grown in the presence of IFN for 23 hours. The IFN-containing medium was removed and cells were incubated with virus (EMC virus or HAV) for 1 hour. After removing the virus, the cells were left for 24 hrs and viable cells stained with methyl violet. The number of viable cells was determined by measuring the optical density of each well. Duplicates of six five fold dilutions of IFN were included in each assay and each assay was repeated at least four times. Antiviral activity of each subtype was compared to lymphoblastoid IFN (WELLFERON[™]) of known activity (10^8 IU/ml).

RESULTS

a) Antiviral Activity

The antiviral effects of some of the different IFN subtypes in HuH7 (liver), A549 (lung) and SHSY (neuroblastoma) cells challenged with EMC virus are shown in Figure 1, which gives the ED50 (dose of IFN causing 50% inhibition of viral replication) for all different subtypes.

The efficacy of the different subtypes was similar in

another liver cell line (HepG2) (data not shown) and a similar trend was seen when HuH7 cells were challenged with another virus (hepatitis A virus). There was a marked difference in the relative efficacies of the different subtypes between the cell lines: in liver cell lines IFN- α_8 , - α_5 and - α_2 were very potent whilst IFN- α_{17} had relatively little antiviral activity. In lung carcinoma cell lines IFN- α_8 , - α_{17} , - α_{10} , - α_5 and - α_{14} were the most potent subtypes tested. IFN- α_1 had very poor activity.

When the anti-viral effects of the different subtypes were analysed using the interferon resistant cell line 11,1, only IFN- α_8 inhibited the cytopathic effects, indicating that this type of subtype has unique properties not shared by the other IFN- α subtypes.

DISCUSSION

1. Between cell lines there are differences in antiviral activity between different subtypes; therefore it may be necessary to use specific subtypes for treatment of infection of each cell type in clinical practice: infections of liver and lung may require different IFN subtypes.

2. IFN- α_2 , IFN- α_8 and - α_5 are active against both liver and lung cell lines; therefore they may induce most side effects in patients - should use a cell specific subtype to reduce side effects (eg α_8 for hepatocytes).

3. However, in view of cells' response to viral infection, i.e production of IFN- α_8 , this subtype may be the subtype of choice generally.

4. In view of the specific activity of IFN- α_8 in mutated cell lines, this subtype may have additional desirable properties and may be effective when other subtypes are inactive.

CLAIMS


1. The use of a single interferon- α (IFN- α) subtype in the preparation of a medicament for preventing or
5 treating viral infections of a particular organ or cell type.
2. The use as claimed in claim 1, wherein the subtype has potent antiviral activity for that particular cell
10 type and relatively low activity in respect of other cell types.
3. The use of IFN- α_{10} and/or IFN- α_{17} in the preparation of a medicament for preventing or treating viral
15 infections of the lung.
4. The use of IFN- α_8 in the preparation of a medicament for preventing or treating viral infections of a
20 particular organ or cell type.
5. The use as claimed in claim 4, wherein the organ is the liver.
6. The use as claimed in claim 5, wherein the viral
25 infection is caused by Hepatitis virus.
7. A method of preventing or treating viral infections of a particular organ or cell type, the method comprising administering an effective antiviral amount of a single
30 interferon- α (IFN- α) subtype.
8. A method as claimed in claim 6, wherein the interferon- α (IFN- α) subtype is IFN- α_8 .

9. A method as claimed in claim 7, wherein the organ is the liver.


10. A method as claimed in claim 9, wherein the viral
5 infection is caused by Hepatitis virus.

1 / 1


ED 50 OF DIFFERENT INTERFERON SUBTYPES IN HuH7 CELLS



ED 50 OF DIFFERENT SUBTYPES IN SHSY CELLS



ED 50 OF DIFFERENT INTERFERON SUBTYPES IN A549 CELLS



INTERNATIONAL SEARCH REPORT

Int. onal Application No

PCT/GB 95/00488

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K38/21

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BUNDESVERBAND DER PHARMAZEUTISCHE INDUSTRIE BV. 'Rote Liste 1993' 1993, EDITIO CANTOR, AULENDORF DE see abstracts 50028 and 50032 ---	1,2,7
X	ANTIVIRAL RESEARCH, vol. 22, no. 2-3, 1993 AMSTERDAM, pages 121-129, SPERBER, S.J. ET AL 'Anti-rhinoviral activity of recombinant and hybrid species of interferon alpha' see page 125 - page 126 --- -/--	1-4,7,8

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- 'A' document defining the general state of the art which is not considered to be of particular relevance
- 'E' earlier document but published on or after the international filing date
- 'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- 'O' document referring to an oral disclosure, use, exhibition or other means
- 'P' document published prior to the international filing date but later than the priority date claimed

- 'T' later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- 'X' document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- 'Y' document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- '&' document member of the same patent family

Date of the actual completion of the international search

31 May 1995

Date of mailing of the international search report

20.06.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Fernandez y Branas, F

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 95/00488

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,93 16107 (INTERFERON SCIENCES) 19 August 1993 see page 25 see page 35 - page 36 see page 51; table 11 see page 111; claim 1 ---	1-10
X,P	CLINICAL SCIENCE, vol. 88,no. 2, February 1995 GLASGOW UK, page 13p FOSTER GR ET AL 'Induction of, and response to, different interferon alpha subtypes in human cell lines' see abstract n45 ---	1-10
X	VIROLOGY, vol. 130, 1983 NEW YORK, pages 273-280, TAMAR GOREN ET AL 'High and low potency interferon-alpha subtypes induce (2'-5') oligoadenylate synthetase with similar efficiency' see the whole document ---	1,2,7
X	JOURNAL OF INTERFERON RESEARCH, vol. special issue, January 1991 CHICAGO, pages 185-194, FINTER N.B. 'Why are there so many subtypes of Alpha-interferons?' see the whole document ---	1,2,4,7, 8
A	EP,A,0 043 980 (GENENTECH INC) 1982 see the whole document ---	1-10
A	JOURNAL OF INTERFERON RESEARCH, vol. 9, 1989 CHICAGO, pages 97-114, FISH E.N. ET AL 'The role of three domains in the biological activity of human Interferon-alpha' see the whole document ---	1-10
A	WO,A,84 03105 (SECHER D.S.) 1984 see page 3 ---	1-10
A	REYNOLDS, J.E.F. 'Martindale, The extra Pharmacopoeia' 1989 , THE PHARMACEUTICAL PRESS , LONDON see page 696 - page 699 -----	1-10

INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB95/00488

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 7-10 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

In ternational Application No

PCT/GB 95/00488

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9316107	19-08-93	AU-B- 3660093 CA-A- 2129533 EP-A- 0625991 JP-T- 7503851	03-09-93 19-08-93 30-11-94 27-04-95
EP-A-43980	20-01-82	AT-B- 394207 AT-B- 394208 AU-B- 552004 AU-A- 7246281 CH-A- 657141 CH-A- 651308 DE-A- 3125706 DE-A- 3177288 EP-A,B 0211148 FR-A,B 2486098 GB-A,B 2079291 JP-C- 1831769 JP-A- 57079897 JP-B- 63061960 JP-C- 1780723 JP-A- 60227694 JP-B- 63061920 JP-C- 1778848 JP-A- 60221093 JP-B- 63063198 JP-C- 1778849 JP-A- 60221094 JP-B- 63063199 LU-A- 83461 NL-A- 8103151 OA-A- 6845 SE-B- 465223 SE-A- 8104093 SU-A- 1414319	25-02-92 25-02-92 22-05-86 07-01-82 15-08-86 13-09-85 15-04-82 01-10-92 25-02-87 08-01-82 20-01-82 29-03-94 19-05-82 30-11-88 13-08-93 12-11-85 30-11-88 13-08-93 05-11-85 06-12-88 13-08-93 05-11-85 06-12-88 06-04-83 01-02-82 28-02-83 12-08-91 02-01-82 30-07-88
WO-A-8403105	16-08-84	EP-A- 0134803 GB-A,B 2146351 JP-T- 60500714	27-03-85 17-04-85 16-05-85